

## 12.1

## DOUBLE INTEGRALS OVER RECTANGLES

**EXAMPLE A** The contour map in Figure 1 shows the snowfall, in inches, that fell on the state of Colorado on December 24, 1982. (The state is in the shape of a rectangle that measures 388 mi west to east and 276 mi south to north.) Use the contour map to estimate the average snowfall for Colorado as a whole on December 24. The **average value** of a function  $f$  of two variables defined on a rectangle  $R$  is

$$f_{\text{ave}} = \frac{1}{A(R)} \iint_R f(x, y) dA$$

where  $A(R)$  is the area of  $R$ .




FIGURE 1

**SOLUTION** Let's place the origin at the southwest corner of the state. Then  $0 \leq x \leq 388$ ,  $0 \leq y \leq 276$ , and  $f(x, y)$  is the snowfall, in inches, at a location  $x$  miles to the east and  $y$  miles to the north of the origin. If  $R$  is the rectangle that represents Colorado, then the average snowfall for the state on December 24 was

$$f_{\text{ave}} = \frac{1}{A(R)} \iint_R f(x, y) dA$$

where  $A(R) = 388 \cdot 276$ . To estimate the value of this double integral let's use the Midpoint Rule with  $m = n = 4$ . In other words, we divide  $R$  into 16 subrectangles of equal size, as in Figure 2. The area of each subrectangle is

$$\Delta A = \frac{1}{16}(388)(276) = 6693 \text{ mi}^2$$




FIGURE 2

Using the contour map to estimate the value of  $f$  at the center of each subrectangle, we get

$$\begin{aligned} \iint_R f(x, y) dA &\approx \sum_{i=1}^4 \sum_{j=1}^4 f(\bar{x}_i, \bar{y}_j) \Delta A \\ &\approx \Delta A [0.4 + 1.2 + 1.8 + 3.9 + 0 + 3.9 + 4.0 + 6.5 \\ &\quad + 0.1 + 6.1 + 16.5 + 8.8 + 1.8 + 8.0 + 16.2 + 9.4] \\ &= (6693)(88.6) \end{aligned}$$

$$\text{Therefore } f_{\text{ave}} \approx \frac{(6693)(88.6)}{(388)(276)} \approx 5.5$$

On December 24, 1982, Colorado received an average of approximately  $5\frac{1}{2}$  inches of snow. ■