

### 4.3


## DISCOVERY PROJECT: AREA FUNCTIONS

This project can be completed anytime after you have studied Section 4.3 in the textbook.

1. (a) Draw the line  $y = 2t + 1$  and use geometry to find the area under this line, above the  $t$ -axis, and between the vertical lines  $t = 1$  and  $t = 3$ .  
 (b) If  $x > 1$ , let  $A(x)$  be the area of the region that lies under the line  $y = 2t + 1$  between  $t = 1$  and  $t = x$ . Sketch this region and use geometry to find an expression for  $A(x)$ .  
 (c) Differentiate the area function  $A(x)$ . What do you notice?
2. (a) If  $0 \leq x \leq \pi$ , let  $A(x) = \int_0^x \sin t \, dt$ .  $A(x)$  represents the area of a region. Sketch that region.  
 (b) Use the Evaluation Theorem to find an expression for  $A(x)$ .  
 (c) Find  $A'(x)$ . What do you notice?  
 (d) If  $x$  is any number between 0 and  $\pi$  and  $h$  is a small positive number, then  $A(x + h) - A(x)$  represents the area of a region. Describe and sketch the region.  
 (e) Draw a rectangle that approximates the region in part (d). By comparing the areas of these two regions, show that

$$\frac{A(x + h) - A(x)}{h} \approx \sin x$$

- (f) Use part (e) to give an intuitive explanation for the result of part (c).

-  3. (a) Draw the graph of the function  $f(x) = \cos(x^2)$  in the viewing rectangle  $[0, 2]$  by  $[-1.25, 1.25]$ .  
 (b) If we define a new function  $g$  by

$$g(x) = \int_0^x \cos(t^2) \, dt$$

then  $g(x)$  is the area under the graph of  $f$  from 0 to  $x$  [until  $f(x)$  becomes negative, at which point  $g(x)$  becomes a difference of areas]. Use part (a) to determine the value of  $x$  at which  $g(x)$  starts to decrease. [Unlike the integral in Problem 2, it is impossible to evaluate the integral defining  $g$  to obtain an explicit expression for  $g(x)$ .]

- (c) Use the integration command on your calculator or computer to estimate  $g(0.2)$ ,  $g(0.4)$ ,  $g(0.6)$ ,  $\dots$ ,  $g(1.8)$ ,  $g(2)$ . Then use these values to sketch a graph of  $g$ .  
 (d) Use your graph of  $g$  from part (c) to sketch the graph of  $g'$  using the interpretation of  $g'(x)$  as the slope of a tangent line. How does the graph of  $g'$  compare with the graph of  $f$ ?
4. Suppose  $f$  is a continuous function on the interval  $[a, b]$  and we define a new function  $g$  by the equation

$$g(x) = \int_a^x f(t) \, dt$$

Based on your results in Problems 1–3, conjecture an expression for  $g'(x)$ .