

3.4 EXPONENTIAL GROWTH AND DECAY**A** [Click here for answers.](#)

1. A bacteria culture starts with 4000 bacteria and the population triples every half-hour.
 - (a) Find an expression for the number of bacteria after t hours.
 - (b) Find the number of bacteria after 20 min.
 - (c) When will the population reach 20,000?
2. A bacteria culture grows with constant relative growth rate. The count was 400 after 2 hours and 25,600 after 6 hours.
 - (a) What was the initial population of the culture?
 - (b) Find an expression for the population after t hours.
 - (c) In what period of time does the population double?
 - (d) When will the population reach 100,000?
3. Polonium-210 has a half-life of 140 days.
 - (a) If a sample has a mass of 200 mg, find a formula for the mass that remains after t days.
 - (b) Find the mass after 100 days.
 - (c) When will the mass be reduced to 10 mg?
 - (d) Sketch the graph of the mass function.

S [Click here for solutions.](#)


4. Polonium-214 has a very short half-life of 1.4×10^{-4} s.
 - (a) If a sample has a mass of 50 mg, find a formula for the mass that remains after t seconds.
 - (b) Find the mass that remains after a hundredth of a second.
 - (c) How long would it take for the mass to decay to 40 mg?
5. On a hot day a thermometer is taken outside from an air-conditioned room where the temperature is 21°C . After one minute it reads 27°C and after 2 minutes it reads 30°C .
 - (a) What is the outdoor temperature?
 - (b) Sketch the graph of the temperature function.

3.4

ANSWERS


E [Click here for exercises.](#)

1. (a) $y(t) = 4000 \cdot 9^t$
 (b) 8320
 (c) ≈ 44 min
2. (a) 50
 (b) $y(t) = 50 \cdot 8^{t/2}$
 (c) 40 min.
 (d) ≈ 7.3 h
3. (a) $y(t) = 200 \cdot 2^{-t/140}$
 (b) ≈ 121.9 mg
 (c) ≈ 605 days
 (d)



S [Click here for solutions.](#)

4. (a) $y(t) = 50 \cdot 2^{-t/0.00014}$
 (b) $\approx 1.57 \times 10^{-20}$ mg
 (c) $\approx 4.5 \times 10^{-5}$ s
5. (a) 33°
 (b)



3.4 SOLUTIONS



1. (a) By (2), $y(t) = y(0)e^{kt} = 4000e^{kt} \Rightarrow$
 $y\left(\frac{1}{2}\right) = 4000e^{k/2} = 12,000 \Rightarrow e^{k/2} = 3 \Rightarrow$
 $k/2 = \ln 3 \Rightarrow k = 2 \ln 3$, so
 $y(t) = 4000e^{(2 \ln 3)t} = 4000 \cdot 9^t$.

(b) $y\left(\frac{1}{3}\right) = 4000 \cdot 9^{1/3} \approx 8320$

(c) $4000 \cdot 9^t = 20,000 \Rightarrow 9^t = 5 \Rightarrow t \ln 9 = \ln 5$
 $\Rightarrow t = (\ln 5) / (\ln 9) \approx 0.73 \text{ h} \approx 44 \text{ min}$

2. (a) $y(t) = y(0)e^{kt} \Rightarrow y(2) = y(0)e^{2k} = 400$,
 $y(6) = y(0)e^{6k} = 25,600$. Dividing these equations,
 we get $e^{6k}/e^{2k} = 25,600/400 \Rightarrow e^{4k} = 64 \Rightarrow$
 $4k = \ln 64 = 6 \ln 2 \Rightarrow k = \frac{3}{2} \ln 2 = \frac{1}{2} \ln 8$. Thus,
 $y(0) = 400/e^{2k} = 400/e^{\ln 8} = \frac{400}{8} = 50$.

(b) $y(t) = y(0)e^{kt} = 50e^{(\ln 8)t/2}$ or $y = 50 \cdot 8^{t/2}$


(c) $y(t) = 50e^{(3 \ln 2)t/2} = 100 \Leftrightarrow e^{(3 \ln 2)t/2} = 2 \Leftrightarrow$
 $(3 \ln 2)t/2 = \ln 2 \Leftrightarrow t = 2/3 \text{ h} = 40 \text{ min}$

(d) $50e^{(\ln 8)t/2} = 100,000 \Leftrightarrow e^{(\ln 8)t/2} = 2000$
 $\Leftrightarrow (\ln 8)t/2 = \ln 2000 \Leftrightarrow$
 $t = (2 \ln 2000) / \ln 8 \approx 7.3 \text{ h}$.

3. (a) The mass remaining after t days is
 $y(t) = y(0)e^{kt} = 200e^{kt}$. Since the half-life is
 140 days, $y(140) = 200e^{140k} = 100 \Rightarrow e^{140k} = \frac{1}{2}$
 $\Rightarrow 140k = \ln \frac{1}{2} \Rightarrow k = -(\ln 2)/140$, so
 $y(t) = 200e^{-(\ln 2)t/140} = 200 \cdot 2^{-t/140}$.

(b) $y(100) = 200 \cdot 2^{-100/140} \approx 121.9 \text{ mg}$

(c) $200e^{-(\ln 2)t/140} = 10 \Leftrightarrow$
 $-\ln 2 \frac{t}{140} = \ln \frac{1}{20} = -\ln 20 \Leftrightarrow$
 $t = 140 \frac{\ln 20}{\ln 2} \approx 605 \text{ days}$



4. (a) If $y(t)$ is the mass remaining after t days,
 then $y(t) = y(0)e^{kt} = 50e^{kt}$.
 $y(0.00014) = 50e^{0.00014k} = 25 \Rightarrow$
 $e^{0.00014k} = \frac{1}{2} \Rightarrow k = -(\ln 2)/0.00014 \Rightarrow$
 $y(t) = 50e^{-(\ln 2)t/0.00014} = 50 \cdot 2^{-t/0.00014}$

(b) $y(0.01) = 50 \cdot 2^{-0.01/0.00014} \approx 1.57 \times 10^{-20} \text{ mg}$

(c) $50e^{-(\ln 2)t/0.00014} = 40 \Rightarrow$
 $-(\ln 2)t/0.00014 = \ln 0.8 \Rightarrow$
 $t = -0.00014 \frac{\ln 0.8}{\ln 2} \approx 4.5 \times 10^{-5} \text{ s}$

5. (a) Let $y(t)$ = temperature after t minutes. Newton's Law
 of Cooling implies that $\frac{dy}{dt} = k(y - a)$ where a is the
 surrounding temperature. Let $u(t) = y(t) - a$. Then
 $\frac{du}{dt} = ku$, so $u(t) = u(0)e^{kt} = (21 - a)e^{kt}$ and
 so $y(t) = a + (21 - a)e^{kt}$. Using the
 knowledge of $y(t)$ at $t = 1$ and $t = 2$ we
 have: $27 = y(1) = a + (21 - a)e^k$ and
 $30 = y(2) = a + (21 - a)e^{2k}$. Rearranged,
 these become $27 - a = (21 - a)e^k$ and
 $30 - a = (21 - a)e^{2k}$. To determine a we must
 eliminate k . To do so, we divide the square of
 the first equation by the second and get
 $\frac{(27 - a)^2}{30 - a} = \frac{(21 - a)^2 e^{2k}}{(21 - a) e^{2k}} \Rightarrow \frac{(27 - a)^2}{30 - a} = 21 - a$
 $\Rightarrow (27 - a)^2 = (30 - a)(21 - a) \Leftrightarrow$
 $729 - 54a + a^2 = 630 - 51a + a^2 \Rightarrow -3a = -99$
 $\Rightarrow a = 33$. Hence, the outdoor temperature is 33° .

