

12.8 CHANGE OF VARIABLES IN MULTIPLE INTEGRALS**A** Click here for answers.**1–6** ■ Find the Jacobian of the transformation.

1. $x = u - 2v, \quad y = 2u - v$

2. $x = u - v^2, \quad y = u + v^2$

3. $x = e^{2u} \cos v, \quad y = e^{2u} \sin v$

4. $x = se^t, \quad y = se^{-t}$

5. $x = u + v + w, \quad y = u + v - w, \quad z = u - v + w$

6. $x = 2u, \quad y = 3v^2, \quad z = 4w^3$

7–8 ■ Find the image of the set S under the given transformation.

7. $S = \{(u, v) \mid 0 \leq u \leq 2, 0 \leq v \leq 1\};$
 $x = u - 2v, \quad y = 2u - v$

8. $S = \{(u, v) \mid 0 \leq u \leq 1, u \leq v \leq 1\};$
 $x = u^2, \quad y = v$

S Click here for solutions.**9–10** ■ Use the given transformation to evaluate the integral.

9. $\iint_R (3x + 4y) \, dA$, where R is the region bounded by the lines $y = x$, $y = x - 2$, $y = -2x$, and $y = 3 - 2x$;
 $x = \frac{1}{3}(u + v), \quad y = \frac{1}{3}(v - 2u)$

10. $\iint_R (x + y) \, dA$, where R is the square with vertices $(0, 0)$, $(2, 3)$, $(5, 1)$, and $(3, -2)$; $x = 2u + 3v, \quad y = 3u - 2v$

11–12 ■ Evaluate the integral by making an appropriate change of variables.

11. $\iint_R xy \, dA$, where R is the region bounded by the lines $2x - y = 1$, $2x - y = -3$, $3x + y = 1$, and $3x + y = -2$

12. $\iint_R \frac{x + 2y}{\cos(x - y)} \, dA$, where R is the parallelogram bounded by the lines $y = x$, $y = x - 1$, $x + 2y = 0$, and $x + 2y = 2$

12.8**ANSWERS**[E Click here for exercises.](#)[S Click here for solutions.](#)

1. 3
2. $4v$
3. $2e^{4u}$
4. $-2s$
5. -4
6. $144vw^2$
7. The parallelogram bounded by the lines $y = 2x$, $y = 2x + 3$,
 $x = 2y$, $x = 2y - 6$
8. The figure bounded by the lines $x = 0$, $y = 1$ and the
parabola $x = y^2$
9. $\frac{11}{3}$
10. 39
11. $-\frac{66}{125}$
12. $\frac{2}{3} \ln(\sec 1 + \tan 1)$

12.8 SOLUTIONS

Click here for exercises.

$$1. \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \partial x / \partial u & \partial x / \partial v \\ \partial y / \partial u & \partial y / \partial v \end{vmatrix} = \begin{vmatrix} 1 & -2 \\ 2 & -1 \end{vmatrix} \\ = 1(-1) - 2(-2) = 3$$

$$2. \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \partial x / \partial u & \partial x / \partial v \\ \partial y / \partial u & \partial y / \partial v \end{vmatrix} = \begin{vmatrix} 1 & -2v \\ 1 & 2v \end{vmatrix} \\ = 2v - (-2v) = 4v$$


$$3. \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \partial x / \partial u & \partial x / \partial v \\ \partial y / \partial u & \partial y / \partial v \end{vmatrix} = \begin{vmatrix} 2e^{2u} \cos v & -e^{2u} \sin v \\ 2e^{2u} \sin v & e^{2u} \cos v \end{vmatrix} \\ = 2e^{4u} (\cos^2 v + \sin^2 v) = 2e^{4u}$$

$$4. \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \partial x / \partial u & \partial x / \partial v \\ \partial y / \partial u & \partial y / \partial v \end{vmatrix} = \begin{vmatrix} e^t & se^t \\ e^{-t} & -se^{-t} \end{vmatrix} \\ = -s - s = -2s$$


$$5. \frac{\partial(x, y, z)}{\partial(u, v, w)} = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 1 & -1 \\ 1 & -1 & 1 \end{vmatrix} \\ = 1(1-1) - 1(1+1) + 1(-1-1) = -4$$

$$6. \frac{\partial(x, y, z)}{\partial(u, v, w)} = \begin{vmatrix} 2 & 0 & 0 \\ 0 & 6v & 0 \\ 0 & 0 & 12w^2 \end{vmatrix} \\ = (2)(6v)(12w^2) = 144vw^2$$

7. S_1 : $v = 0, 0 \leq u \leq 2$, so $x = u, y = 2u$ and $y = 2x$.
 S_2 : $u = 2, 0 \leq v \leq 1$, so $x = 2 - 2v, y = 4 - v$ and $x = 2y - 6$.
 S_3 : $v = 1, 0 \leq u \leq 2$, so $x = u - 2, y = 2u - 1$ and $y = 2x + 3$.
 S_4 : $u = 0, 0 \leq v \leq 1$, so $x = -2v, y = -v$ and $2y = x$.



8. S_1 : $u = v, 0 \leq u \leq 1$, so $x = u^2, y = u$, and $x = y^2$.
 S_2 : $v = 1, 0 \leq u \leq 1$, so $x = u^2, y = 1$.
 S_3 : $u = 0, 0 \leq v \leq 1$, so $x = 0, y = v$.



$$9. \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{1}{3} & \frac{1}{3} \\ -\frac{2}{3} & \frac{1}{3} \end{vmatrix} = \frac{1}{3} \text{ and}$$

$3x + 4y = (u + v) + \frac{4}{3}(v - 2u) = \frac{1}{3}(7v - 5u)$. To find the region S in the uv -plane that corresponds to R we first find the corresponding boundary under the given transformation: The line $y = x$ is the image of

$$\frac{1}{3}(v - 2u) = \frac{1}{3}(u + v) \text{ or } u = 0, y = x - 2 \Rightarrow$$

$$\frac{1}{3}(v - 2u) = \frac{1}{3}(u + v) - 2 \text{ or } u = 2, y = -2x \Rightarrow$$

$$\frac{1}{3}(v - 2u) = -\frac{2}{3}(u + v) \text{ or } v = 0, \text{ and } y = 3 - 2x \Rightarrow$$

$$\frac{1}{3}(v - 2u) = 3 - \frac{2}{3}(u + v) \text{ or } v = 3. \text{ Thus } S \text{ is the}$$

rectangle $[0, 2] \times [0, 3]$ in the uv -plane and

$$\iint_R (3x + 4y) dA = \iint_S \frac{1}{3}(7v - 5u) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv \\ = \int_0^3 \int_0^2 \frac{1}{3}(7v - 5u) \left(\frac{1}{3} \right) du dv \\ = \frac{1}{9} \int_0^3 (14v - 10) dv = \frac{1}{9}(33) = \frac{11}{3}$$

$$10. \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} 2 & 3 \\ 3 & -2 \end{vmatrix} = -13, x + y = 5u + v \text{ and since}$$

$u = \frac{2x + 3y}{13}$ and $v = \frac{3x - 2y}{13}$, R is the image of the square with vertices $(0, 0), (1, 0), (1, 1), (0, 1)$. Thus

$$\iint_R (x + y) dA = \int_0^1 \int_0^1 (5u + v) |-13| du dv \\ = 13 \int_0^1 \left(\frac{5}{2} + v \right) dv = 13(3) = 39$$

11. Letting $u = 2x - y$ and $v = 3x + y$, we have $x = \frac{1}{5}(u + v)$,

$$y = \frac{1}{5}(2v - 3u). \text{ Then } \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{1}{5} & \frac{1}{5} \\ -\frac{3}{5} & \frac{2}{5} \end{vmatrix} = \frac{1}{5} \text{ and}$$

$$\iint_R xy dA = \int_{-2}^1 \int_{-3}^1 \frac{(u + v)(2v - 3u)}{25} \left(\frac{1}{5} \right) du dv \\ = \frac{1}{125} \int_{-2}^1 \int_{-3}^1 (2v^2 - uv - 3u^2) du dv \\ = \frac{1}{125} \int_{-2}^1 (8v^2 + 4v - 28) dv = -\frac{66}{125}$$

12. Let $u = x - y, v = x + 2y$, so $y = \frac{1}{3}(v - u)$ and

$$x = \frac{1}{3}(2u + v). \text{ Then } \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{2}{3} & \frac{1}{3} \\ -\frac{1}{3} & \frac{1}{3} \end{vmatrix} = \frac{1}{3} \text{ and}$$

$$\iint_R \frac{x + 2y}{\cos(x - y)} dA = \frac{1}{3} \int_0^1 \int_0^2 \frac{v}{\cos u} dv du \\ = \frac{2}{3} \int_0^1 \sec u du = \frac{2}{3} [\ln |\sec u + \tan u|]_0^1 \\ = \frac{2}{3} [\ln(\sec 1 + \tan 1) - \ln 1] = \frac{2}{3} \ln(\sec 1 + \tan 1)$$